


Figure 3. Weed susceptibility, application timing and water management regimes for herbicide sequences with Regiment.

	barnyardgrass	watergrass	sprangletop	smallflower umbrella	ricefield bulrush	CA arrowhead	Gregg's arrowhead	ducksalad	redstem	monochoria	
Bolero	+ / R	+ / R	+	+	-	-	-	-	-	-	+ control - no control ± suppression R resistant
Followed by											
Regiment	+ / R	+ / R	-	-	+ / R	+ / R	-	-	-	±	

Bolero fb. Regiment (Permanent Flood)



1.0 to 2.0 lsr (4.0 lb ai/ac)

Application timing


Fb.

1-3 til (15 g ai/ac)

	barnyardgrass	watergrass	sprangletop	smallflower umbrella	ricefield bulrush	CA arrowhead	Gregg's arrowhead	ducksalad	redstem	monochoria
--	---------------	------------	-------------	----------------------	-------------------	--------------	-------------------	-----------	---------	------------

Regiment	+ / R	+ / R	-	-	+ / R	+ / R	-	-	-	±	+ control - no control ± suppression R resistant
Followed by											
Stam or Superwham	+	+	-	+	+	±	-	±	±	±	

Regiment fb. propanil (Pin-point Flood)



Application timing

**NO RESISTANCE TO
PROPANIL**

5 lsr to 1 til (15 g ai/ac)

Fb.

2-3 til (6 lb ai/ac)